

 OPEN HERE

FMC Agro Limited, Rectors Lane, Pentre, Flintshire, CH5 2DH.
Tel: 01244 537370. Email: Flintshire.enquiry@fmc.com
Website: www.fmc-agro.co.uk. Technical Helpline: 01423 205011


Emergency No.s - Medical Incidents Involving Crop Protection

Products: England & Wales: 111, Scotland: 08454 24 24 24, Northern Ireland: Local GP or Pharmacist, Republic of Ireland: 01 837 9964. Healthcare professionals seeking poisons information: www.toxbase.org.

Emergency No.s - Transportation, Warehousing & After Sale Incidents:

CHEMTREC help centre Dialling from the UK and NI: 0870 820 0418, Dialling from Ireland: 01 901 4670

Manufactured in the E.U.



Very toxic to aquatic life with long lasting effects.

• Wear protective gloves/ protective clothing. IF ON SKIN: Wash with plenty of soap and water. • If skin irritation or rash occurs: Get medical advice/attention • Wash contaminated clothing before reuse • Collect spillage. • Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste • Do not contaminate water with the product or its container (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

To avoid risks to human health and the environment, comply with the instructions for use.

F-05264/31908 - UK



CAMEO

Herbicide

MAPP 18990 

A soluble granule containing 500 g/kg tribenuron-methyl, a sulfonylurea, for broad-leaved weed control in wheat, oats, barley, durum wheat, triticale, winter rye and green cover on land not being used for crop production.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

Contents: 120 g e



IMPORTANT INFORMATION

FOR USE ONLY AS A PROFESSIONAL HERBICIDE

Maximum individual dose:	30 g/ha
Maximum number of treatments:	One per crop
Crops:	wheat, barley, oats, rye (winter), triticale, durum wheat and green cover on land not being used for crop production
Latest time of application:	GS39

Other Specific Restrictions

This product must not be applied at rates greater than 10 g product/ha before the end of February in the year of harvest.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR PLANT PROTECTION PRODUCTS.

READ DIRECTIONS FOR USE ON ATTACHED LEAFLET FOR PROFESSIONAL USE ONLY



The Voluntary Initiative
This label is compliant with the CPA Voluntary Initiative Guidance



To access the Safety Data Sheet for this product scan the code

Batch N°:

FMC Agro Limited, Rectors Lane, Pentre, Flintshire, CH5 2DH.
Tel: 01244 537370. Email: Flintshire.enquiry@fmc.com
Website: www.fmc-agro.co.uk. Technical Helpline: 01423 205011



Emergency No.s - Medical Incidents Involving Crop Protection

Products: England & Wales: 111, Scotland: 08454 24 24 24, Northern Ireland: Local GP or Pharmacist, Republic of Ireland: 01 837 9964. Healthcare professionals seeking poisons information: www.toxbase.org.

Emergency No.s - Transportation, Warehousing & After Sale Incidents:

CHEMTREC help centre Dialling from the UK and NI: 0870 820 0418, Dialling from Ireland: 01 901 4670

Manufactured in the E.U.

	Herbicide contains 500 g/kg tribenuron-methyl	
	WARNING May cause an allergic skin reaction	

Very toxic to aquatic life with long lasting effects.

- Wear protective gloves/ protective clothing. IF ON SKIN: Wash with plenty of soap and water. • If skin irritation or rash occurs: Get medical advice/attention • Wash contaminated clothing before reuse
- Collect spillage. • Dispose of contents/container to a licensed hazardous-waste disposal contractor or collection site except for empty clean containers which can be disposed of as non-hazardous waste • Do not contaminate water with the product or its container (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

To avoid risks to human health and the environment, comply with the instructions for use.

F-05264/31908 - UK

FMC

CAMEO

Herbicide

MAPP 18990 

A soluble granule containing 500 g/kg tribenuron-methyl, a sulfonyleurea, for broad-leaved weed control in wheat, oats, barley, durum wheat, triticale, winter rye and green cover on land not being used for crop production.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

Contents: **120 g e**



IMPORTANT INFORMATION

FOR USE ONLY AS A PROFESSIONAL HERBICIDE

Maximum individual dose:	30 g/ha
Maximum number of treatments:	One per crop
Crops:	wheat, barley, oats, rye (winter), triticale, durum wheat and green cover on land not being used for crop production
Latest time of application:	GS39
Other Specific Restrictions	

This product must not be applied at rates greater than 10 g product/ha before the end of February in the year of harvest.

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR PLANT PROTECTION PRODUCTS.

READ DIRECTIONS FOR USE ON ATTACHED LEAFLET FOR PROFESSIONAL USE ONLY


The Voluntary Initiative
This label is compliant with the
CPA Voluntary Initiative Guidance



To access the Safety Data Sheet for this product scan the code

CAMEO

Herbicide

MAPP 18990



A soluble granule containing 500 g/kg tribenuron-methyl, a sulfonylurea, for broad-leaved weed control in wheat, oats, barley, durum wheat, triticale, winter rye and green cover on land not being used for crop production.

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

FMC Agro Limited, Rectors Lane, Pentre, Flintshire, CH5 2DH.

Tel: 01244 537370. Email: Flintshire.enquiry@fmc.com

Website: www.fmc-agro.co.uk. Technical Helpline: 01423 205011

Emergency No.s - Medical Incidents Involving Crop Protection Products:

England & Wales: 111, Scotland: 08454 24 24 24, Northern Ireland: Local GP or Pharmacist, Republic of Ireland: 01 837 9964. Healthcare professionals seeking poisons information: www.toxbase.org.

Emergency No.s - Transportation, Warehousing & After Sale Incidents:

CHEMTREC help centre Dialling from the UK and NI: 0870 820 0418, Dialling from Ireland: 01 901 4670

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when handling the product

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WHEN USING DO NOT EAT, DRINK OR SMOKE

WASH CONCENTRATE from skin or eyes immediately

WASH HANDS AND EXPOSED SKIN before eating and drinking and after work

Environmental Protection

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

Storage and Disposal

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS

KEEP OUT OF REACH OF CHILDREN

WASH OUT CONTAINER THOROUGHLY and dispose of safely

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Restrictions

- CAMEO should not be applied to any crop suffering from stress as a result of drought, waterlogging, low temperatures, pest or disease attack, nutrient or lime deficiency or other factors reducing crop growth.
- Do not use CAMEO on cereal crops undersown with grasses, clover or other legumes or any other broad-leaved crop.
- Due to the high level of activity of the herbicide, special care must be taken to avoid damage by drift onto broad-leaved plants outside the target area, or onto ponds, waterways or ditches. Thorough cleansing of equipment is also very important - see below.
- Contract agents should be consulted before using on crops grown for seed.
- CAMEO should not be applied within 7 days of rolling the crop.
- Do not apply CAMEO more than once to any cereal crop.
- All requirements or restrictions on other product labels must be adhered to when applied to the same crop as CAMEO. Contact your distributor for further information.
- Refer to 'Compatibility' for details on use of CAMEO with other 'ALS inhibiting' herbicides

Weed control

CAMEO contains tribenuron methyl, a sulfonylurea herbicide, which works mainly by contact action and is most effective if applied when the weeds are small and actively growing. Good spray cover of the weeds must be obtained for best results. Susceptible plants cease growth almost immediately after application and symptoms can be seen about two weeks after application. Weed control may be reduced when soil conditions are very dry. The susceptibility rating of weeds in the following table refer to good spray cover and good growing conditions.

Weed Species	Plants up to 2 expanded true leaves	Plants up to 6 expanded true leaves	Plants up to 15 cm across/high
Charlock*	S	S	S
Chickweed, Common*	S	S	S
Dead-nettle, Red*	S	S	MR
Fat-hen	S	MS	MS
Forget-me-not*	S	S	MS
Fumitory*	S	S	R
Hemp-nettle, Common	S	S	-
Mayweed*	S	S	S
Poppy, Common*	S	S	R
Redshank	S	S	MS
Shepherd's-purse*	S	S	MS

S = Susceptible MS = Moderately Susceptible MR = Moderately Resistant R = Resistant

* Controlled by autumn or spring application up to six true leaves

In addition to the above weeds, volunteer beans and oilseed rape will be controlled by a dose of 10g/ha.

Resistance

This product contains tribenuron-methyl, which is an ALS inhibitor, also classified by the Herbicide Resistance Action Committee as 'Group B'.

When herbicides with the same mode of action are used repeatedly over several years in the same field, selection of resistant biotypes can take place. These can propagate and may become dominating. A weed species is considered resistant to a herbicide if it survives a correctly applied treatment at the recommended dose. Development of resistance with a weed species can be avoided or delayed by alternating (or tank mixing) with suitable products having a different mode of action.

A strategy for preventing and managing resistance should be adopted. The Weed Resistance Action Group has produced guidelines and copies are available from the HGCA, CPA, your distributor, crop advisor or product manufacturer.

Crops

Cereals

CAMEO can be used post-emergence on all varieties of wheat, barley, oats, winter rye, durum wheat, triticale and land not being used for crop production at the growth stages given below.

Green cover on land not being used for crop production

CAMEO can be used for broad-leaved weed control on land not being used for crop production. Only cereals may be sown in the same calendar year to set-aside land treated with CAMEO.

Dose

Apply CAMEO at 30 grams/hectare.

When CAMEO is applied before the end of February in the year of harvest then no more than 10g product /ha can be applied.

Timing

CAMEO may be applied from the three-leaf stage up to (GS 13) and including the flag leaf fully emerged stage of growth (GS 39).

Following crops

Only cereals, oilseed rape, field beans or grass may be sown in the same calendar year as harvest of a cereal crop treated with CAMEO. In case of crop failure for any reason, sow only cereals within three months of application of CAMEO. Before sowing, soil should be ploughed and cultivated to a depth of at least 15-cm.

Volume and application

BEFORE USING CAMEO, SPRAYING EQUIPMENT MUST BE CLEAN AND FREE FROM CONTAMINATION WITH OTHER PESTICIDES.

Application should be made in 100-200 litres of water per hectare, using suitable ground equipment to give good spray cover of the weeds. Use a conventional field sprayer at a pressure of 2-3 bars and apply as a MEDIUM spray quality (as defined by BCPC).

When crops are thick or weed growth is dense, use 400 litres of water per hectare. Care should be taken not to overlap spray swaths.

Extreme care must be taken to avoid spray drift on to non-crop plants outside of the target area.

Mixing

Quarter fill the spray tank with clean water, start the agitation and add the required quantity of CAMEO directly to the tank without prior creaming. Continue agitation while topping up the tank and while spraying.

Compatibility

To widen the spectrum of activity and improve the control of some weeds, CAMEO may be applied with other herbicides, provided each product can be applied within the label recommendations for its use. Always add CAMEO to the spray tank first when using in tank-mix and ensure it is fully dispersed before adding the partner product.

The following 'joint application' with other sulfonylurea and 'ALS inhibiting' herbicides may be applied to a crop. Requirements and restrictions on other product labels must be adhered to. 'Joint application' is the use of CAMEO in tank mixture or sequence with one of the products listed below.

Absolute	Exceed SX	Hunter	Spitfire
Atlantis WG	Gala	Lexus Class	Spitfire Solo
Barton WG	Galaxy	Lexus Millenium	Staka SX
Broadway Star	Gartrel	Lexus SX	Starane Gold
Broadway Sunrise	GF-184	Oklar SX	Starane Vantage
Boxer	GF-2070	Oriel 50 SX	Starane XL
Bullion	Hatra	Othello	Unite
Dakota	Headland Sure-fire Gold	Pacifica	
Ductis SX	Hiker	Pursuit*	
Eagle*	Horus	Slalom	

* only cereals as following crop

Additional recommendations may be approved since label printing. Please consult your distributor for latest details on compatibilities.

SPRAY TANK CLEAN-OUT

POOR CLEANOUT PRACTICES AND INSUFFICIENT WATER VOLUMES USED FOR THE RINSE PROCEDURE MAY RESULT IN INADEQUATE REMOVAL OF PRODUCT DEPOSITS. SUBSEQUENT USE OF APPLICATION EQUIPMENT IN THESE CIRCUMSTANCES MAY RESULT IN DAMAGE TO NON-CEREAL CROPS.

TO AVOID SUBSEQUENT DAMAGE TO CROPS OTHER THAN CEREALS, IMMEDIATELY AFTER SPRAYING CAMEO THOROUGHLY CLEAN ALL SPRAY EQUIPMENT, INCLUDING INSIDE AND OUTSIDE OF LID, USING THE FOLLOWING PROCEDURE:

Always start with a clean tank and spray system. Clean spray equipment thoroughly immediately after use.

1. Thoroughly and completely rinse all interior tank surfaces (including lid) with water (use at least 10% of the tank capacity), taking care to remove any visible deposits. Flush pump, filters and boom after removing in-line strainers, nozzle tips and screens (clean these parts separately). Drain the remainder of the rinsate from the tank.

2. Repeat the rinse, flush and drain.

3. Dispose of washings safely. Do not spray onto sensitive crop or land intended for cropping with sensitive crop.

Note: Consult label tank cleanup procedures for all tank mix partners and be sure to use the most rigorous procedure recommended.

WARNING

EXTREME CARE SHOULD BE TAKEN TO AVOID DAMAGE BY DRIFT OF SPRAY ONTO BROAD-LEAVED PLANTS OUTSIDE THE TARGET AREA OR ONTO PONDS, WATERWAYS OR DITCHES. SPRAYING EQUIPMENT SHOULD NOT BE DRAINED OR FLUSHED ONTO LAND PLANTED WITH OR INTENDED FOR PLANTING WITH TREES OR CROPS OTHER THAN CEREALS.

NOTICE TO BUYER

All goods supplied by us are of a high grade and we believe them to be suitable for any purpose for which we expressly supply them, but as we cannot exercise control over their mixing or use, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.

All manufacturers trademarks are duly acknowledged.

